arXiv:1210.1824v1 [math.GT] 5 Oct 2012

EXCEPTIONAL SLOPES ON MANIFOLDS OF SMALL COMPLEXITY

FIONNTAN WOUTER MURRAY ROUKEMA

Abstract. It has been observed that most manifolds in the Callahan-Hildebrand-Weeks census of cusped hyperbolic 3-manifolds are obtained by surgery on the minimally twisted 5-chain link; a full classification of the exceptional surgeries on the 5-chain link has recently been completed. In this article, we provide a complete classification of the sets of exceptional slopes and fillings for all cusped hyperbolic surgeries on the minimally twisted 5-chain link, thereby describing the sets of exceptional slopes and fillings for most hyperbolic manifolds of small complexity and providing supporting evidence for some well-known conjectures.

1. Introduction

The set of exceptional slopes on a boundary component of a cusped hyperbolic manifold has generated a lot of interest in the literature; there are many restrictions on the set of exceptional slopes on a boundary component of a hyperbolic 3-manifold M and its corresponding fillings, for example, no such M has two distinct S^3 fillings [GL1] or more than 10 exceptional slopes [LM]. However, it is still not known if a hyperbolic knot exterior in S^3 can have a reducible filling, or if M can have a pair of exceptional slopes α and β corresponding to a lens space and toroidal space so that the *distance* (minimal number of intersections) between α and β is 4. Conjecturally, neither are possible, see [GAS] and [Lee] respectively. Moreover, it is conjecturally the case that the figure-eight knot exterior is the unique one-cusped hyperbolic manifold with 10 exceptional slopes [G1] and that the figure-eight knot exterior and its sister are the unique one-cusped hyperbolic manifolds having two exceptional slopes at distance 8, see [Kir]. In this article, by classifying the sets of exceptional slopes and the corresponding fillings for all manifolds obtained by surgery on the minimally twisted 5-chain link (see the rightmost link in Figure 1), we provide experimental evidence that supports these conjectures:

Theorem 1. *If M is a cusped hyperbolic manifold obtained by surgery on the minimally twisted 5-chain link then:*

- *M is not the exterior of a knot with a reducible filling;*
- *M does not have a lens space and toroidal filling at distance 4;*
- *If M has 10 exceptional slopes then M is the figure-8 knot exterior;*
- *If M has exceptional slopes at distance 8 then M is either the figure-8 knot exterior or the figure-8 knot exterior sister.*

1.1. The minimally twisted 5-chain link.

A notable collection of chain links is described in [MPR]; they are the figure-8 knot, the Whitehead link, the 3-chain link, the 4-chain link with a half twist, and the minimally twisted 5-chain link. These links are shown in Figure 1. We follow [MPR] and denote these chain links by 1CL, 2CL, 3CL, 4CL and 5CL, and their exteriors by M_1, M_2, M_3, M_4, M_5 respectively.



FIGURE 1. The links 1CL, 2CL, 3CL, 4CL, 5CL in S^3 whose exteriors we denote by M_1, M_2, M_3, M_4, and M_5 respectively.

The significance of this sequence of links comes from the following facts: each M_i is the (or conjecturally the) smallest volume hyperbolic 3-manifold with i cusps, see [Ago] and [Yos]; more than 80% of the cusped hyperbolic 3-manifolds from the Callahan-Hildebrand-Weeks census [CHW] are surgeries on 5CL (personal communication with Nathan Dunfield); 5CL relates to the program of enumerating exceptional pairs at maximal distance, in particular, all knots realising half-integral toroidal surgeries [GL2], many of the knots realising lens space surgeries [Bak], and all cusped hyperbolic 3-manifolds with distinct reducible and toroidal fillings at maximal distance [Kan] are obtained by surgery on 5CL.

It is easy to see that if ∂M_n is equipped with the usual (meridian, longitude) homology basis then a -1 filling on any boundary component of M_n results in M_{n-1}, and, as a result, that any manifold obtained by surgery on $(n-1)$CL is obtained by surgery on nCL. A classification of the exceptional surgeries on 3CL is given in [MP] together with a complete description of the set of exceptional slopes, and corresponding exceptional fillings, on the boundary components of all hyperbolic manifolds obtained by surgery on 3CL. A classification of exceptional surgeries on 5CL is found in [MPR]; in this article we completely describe the sets of exceptional slopes and corresponding exceptional fillings of any manifold obtained by surgery on 5CL. To simplify matters, we only consider the 'new' manifolds obtained by surgery on 5CL; namely we only consider surgeries on 5CL and 4CL that are not obtained by surgery on 3CL. In particular, Theorems 3 and 4 provide explicit descriptions of the sets of exceptional slopes and corresponding fillings for all hyperbolic surgeries of 5CL not obtained by surgery on 3CL, and Theorem 1 will be an easy consequence.

1.2. **Article structure.** We start with Section 2 where we recall the classification from [MPR]. In order to do so, we begin by recalling and introducing terminology in Subsection 2.1. The main results of this article are stated and proved in Section 3 and refer to the tables found in Section 4.

1.3. **Acknowledgements.** The results of this article were obtained as a graduate student at the University of Pisa under the supervision of Carlo Petronio and Bruno Martelli. The current presentation of the results has benefited from discussions with Daniel Matignon and Mark Lackenby, and from email correspondences with Carlo Petronio, Bruno Martelli and Nathan Dunfield.

2. Background Terminology and Results

2.1. **Terminology.** We begin with some general terminology, and we introduce the notion of an "exceptional filling instruction" used in Theorems 3 and 4.

Fix an orientable compact 3-manifold X with ∂X consisting of tori:

- A *slope* on a boundary component τ of X is the isotopy class of a non-trivial unoriented loop on τ;
- A *filling instruction* α for X is a set consisting of either a slope or the empty set for each component of ∂X;
- The *filling* $X(\alpha)$ given by an instruction α is the manifold obtained by attaching one solid torus to ∂X for each (non-empty) slope in α, with the meridian of the solid torus attached to the slope.

We recall that if M is a hyperbolic non-compact finite-volume 3-manifold then $M = \text{int}(X)$ with ∂X consisting of tori, and that $\text{int}(X(\alpha))$ is hyperbolic for all but finitely many α's consisting of one slope and $\varnothing$'s [LM].

- If the interior of X is hyperbolic but the interior of $X(\alpha)$ is not, we say that α is an *exceptional filling instruction* for X and that $X(\alpha)$ is an *exceptional filling* of X;
- We say that an exceptional filling instruction α on a hyperbolic X is *isolated* if $X(\beta)$ is hyperbolic for all β properly contained in α; for such an α we call $X(\alpha)$ an *isolated exceptional filling* of X.

A *surgery* on a link L corresponds to a filling of the exterior of L, that is, to a filling of $M\backslash N(L)$ where $N(L)$ is an open regular neighbourhood of L. By a *surgery instruction* for L we mean a filling instruction on the exterior of L.

We now recall some standard notation used in the description of the set of exceptional slopes on a fixed boundary component of a hyperbolic manifold, see for example [G2]. If X is a hyperbolic 3-manifold with boundary consisting of tori and τ is a fixed boundary component of ∂X then the set of exceptional slopes on τ is denoted by $E_\tau(X)$, and the cardinality of $E_\tau(X)$ by $e_\tau(X)$. To describe $E_\tau(M_5(\alpha))$ we introduce the following definition:

Definition Let α be a filling instruction on a manifold X. We say that α *factors through* a manifold Y if there exists some filling instruction $\beta \subset \alpha$ such that $Y = X(\beta)$.

Note that if α is exceptional for X and factors through a hyperbolic Y with $Y = X(\beta)$, then $\alpha\backslash\beta$ is exceptional for Y.

2.2. Exceptional fillings of M_5.

Our description of the exceptional fillings of $M_5(\gamma)$ will employ the notation now discussed for Seifert manifolds with orientable base surface. Given integers $p_1, \ldots, p_n, q_1, \ldots, q_n$, with $p_i > 1$ and G an orientable surface with $k \geq 0$ boundary components $b_1 \ldots b_k$, we let Σ denote the surface obtained by removing n open discs from G and we denote by $b_{k+1}, \ldots, b_{k+n}$ the n newly introduced boundary circles. We fix an orientation on $\Sigma \times S^1$ and orient $\{\mu_i, \lambda_i\} = \{b_i \times \{*\}, \{*\} \times S^1\}$ so that μ_i, λ_i is a positive basis of $H_1(b_i \times S^1)$ with $b_i \times S^1$ oriented as $\partial(\Sigma \times S^1)$. We denote by $(G, (p_1, q_1), \ldots (p_n, q_n))$, the manifold obtained by performing a Dehn filling on each $b_i \times S^1$ along $p_i\mu_i + q_i\lambda_i$ for $i > k$. In our case, G will be either the disc D, the annulus A, or the sphere S^2.

Given Seifert manifolds X and Y with orientable base surfaces with boundary as described above, and $B \in \mathrm{GL}(2, \mathbb{Z})$ with $\det(B) = -1$, we define $X \bigcup_B Y$ unambiguously to be the quotient manifold $X \bigcup_f Y$ where $f : T \to U$ for T and U arbitrary boundary components of X and Y respectively, and f acting on homology by B with respect to the bases described above. The case $T, U \subset \partial X$, $T \neq U$ is also allowed and we write the quotient manifold as $X\big/_{B}$.

For the purposes of Theorem 3 and 4, we employ a flexible notation for Seifert manifolds. We will formally identify an $\varnothing$ slope with $\frac{0}{0}$ and allow parameters (p_i, q_i) to be arbitrary elements of $\mathbb{Z}^2$.

2.3. **Exceptional surgery instructions on 5CL.**

By ordering the components of 5CL cyclically, surgery instructions on 5CL can be naturally identified with $(\mathbb{Q} \cup \{\varnothing, \infty\})^5$. Given such an instruction α, the symmetry group of $M_5(\alpha)$ has a natural action on the boundary components of $M_5(\alpha)$. This action induces an action on the filling instructions on $M_5(\alpha)$. Among the most significant actions arising we mention those coming from the symmetry group of M_5, see (1)-(3) below, a symmetry of M_4 which may be deduced from the Fenn-Rourke blow-down move on 5CL, see (4), and from the amphichirality of the figure-8 knot $M_5(-1, -2, -2, -2, \varnothing)$, see (5); see [MPR] for full details.

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5) \longmapsto (\alpha_5, \alpha_1, \alpha_2, \alpha_3, \alpha_4) \tag{1}$$

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5) \longmapsto (\alpha_5, \alpha_4, \alpha_3, \alpha_2, \alpha_1) \tag{2}$$

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5) \longmapsto (\alpha_2^{-1}, \alpha_1^{-1}, 1 - \alpha_3, (1 - \alpha_4)^{-1}, 1 - \alpha_5) \tag{3}$$

$$(-1, \alpha_1, \alpha_2, \alpha_3, \alpha_4) \longmapsto (\alpha_1, -1, \alpha_2 - 1, \alpha_3, \alpha_4 + 1) \tag{4}$$

$$(-1, -2, -2, -2, \alpha) \longmapsto (-1, -2, -2, -2, -\alpha - 6) \tag{5}$$

Since the action of the maps (1) and (2) is very easily understandable while that of (3)-(5) is more involved, we introduce the symbol $[\alpha]$ for the equivalence class of a filling instruction α under (1) and (2) only, and the symbol $[\![\alpha]\!]$ for the equivalence class of α under (1)-(5).

For a filling instruction α on M_5 we will often simplify notation by omitting empty slopes but leaving the subscripts on non-empty slopes. For example, $((-1)_1, (-1)_3)$ corresponds to the filling instruction $(\varnothing, \mu_1 - \lambda_1, \varnothing, \mu_3 - \lambda_3, \varnothing)$ with (μ_i, λ_i) the (meridian, longitude) basis of the homology of the i^{th} cusp. Note that for any $\frac{p}{q} \in \mathbb{Q} \cup \{\infty\}$ and $i \neq j$ one has $[((\frac{p}{q})_i)] = [((\frac{p}{q})_j)]$, so the fillings $M_5(\frac{p}{q})$ are defined without ambiguity. Our convention will be that a filling instruction $(\frac{p}{q}, \frac{r}{s}, \frac{u}{v}, \frac{x}{y})$ on M_5 with four non-empty slopes and no subscripts represents $(\frac{p}{q}, \frac{r}{s}, \frac{u}{v}, \frac{x}{y}, \varnothing)$. We now state the main result of [MPR].

Theorem 2. *Every exceptional filling instruction on* M_5 *is equivalent up to a composition of the symmetries* (1)-(5) *to a filling instruction containing one of:*

$$\infty, (-1,-2,-2,-1), (-1,-2,-3,-2,-4), (-1,-2,-2,-3,-5),$$
$$(-1,-3,-2,-2,-3), \left(-2,-\tfrac{1}{2},3,3,-\tfrac{1}{2}\right), (-2,-2,-2,-2,-2).$$

Descriptions of the corresponding exceptional fillings may be found in [MPR]. We remark that the second, third, and fourth filling instructions factor through M_3, and we assign the names m_1, m_2, and m_3 to the manifolds $M_5(-1,-3,-2,-2,-3)$, $M_5\left(-2,-\tfrac{1}{2},3,3,-\tfrac{1}{2}\right)$, and $M_5(-2,-2,-2,-2,-2)$ respectively.

3. Main Results

We now precisely describe the classification of exceptional slopes of every hyperbolic surgery on 5CL by describing $E_\tau(M_5(\alpha))$ for every α not factoring through M_3. We split the result according to whether or not α factors through M_4.

Theorem 3. *Let* γ *be a hyperbolic filling instruction on* M_5 *containing at least one* $\varnothing$ *and not factoring through* M_4*, and let* τ *be a boundary component of* $\partial M_5(\gamma)$*; then either* $e_\tau(M_5(\gamma)) = 3$ *and, with respect to the basis induced from* M_5*, we have* $E_\tau(M_5(\gamma)) = \{0,1,\infty\}$ *and*

$$M_5(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f},\tfrac{g}{h})(\infty) = \big(D,(a,-b),(c,d)\big) \bigcup_{\begin{pmatrix}0&1\\1&0\end{pmatrix}} \big(D,(e,f),(g,-h)\big), \tag{6}$$

$$M_5(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f},\tfrac{g}{h})(1) = \big(D,(a-b,b),(e,f)\big) \bigcup_{\begin{pmatrix}0&1\\1&0\end{pmatrix}} \big(D,(e-f,f),(c,d)\big), \tag{7}$$

$$M_5(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f},\tfrac{g}{h})(0) = \big(D,(b,b-a),(h,-g)\big) \bigcup_{\begin{pmatrix}0&1\\1&0\end{pmatrix}} \big(D,(c-d,c),(e-f,f)\big), \tag{8}$$

or $4 \leqslant e_\tau(M_5(\gamma)) \leqslant 5$ *and*

- $M_5(\gamma) \cong M_5(\alpha)$ *for some* α *in Tables* 1-4 *in Section 4, and with respect to the basis induced from* M_5 *we have*

  $$E_\tau(M_5(\alpha)) = \begin{cases} \{\beta_1,\beta_2,0,1,\infty\} & \textit{if } \alpha \textit{ is found in Table } 1; \\ \{\beta,0,1,\infty\} & \textit{otherwise}, \end{cases}$$

  where β, β_1, β_2 *are found in Tables* 1-4.
- *Identities* (6)-(8) *hold and* $M_5(\alpha)(\beta_i)$, $M_5(\alpha)(\beta)$ *are shown in Tables* 1-4.

Ordering the components of M_4 cyclically, identifying filling instructions on M_4 with $(\mathbb{Q}\cup\{\varnothing,\infty\})^4$ and adopting the convention that a filling instruction $(\frac{p}{q},\frac{r}{s},\frac{u}{v})$ on M_4 with three slopes and no subscripts represents $(\frac{p}{q},\frac{r}{s},\frac{u}{v},\varnothing)$, the corresponding result for M_4 is:

Theorem 4. *Let γ be a hyperbolic filling instruction on M_4 containing at least one $\varnothing$ and not factoring through M_3, and let τ a boundary component of $\partial M_4(\gamma)$; then either $e_\tau(M_4(\gamma))=4$ and, with respect to the basis induced from M_4, we have $E_\tau(M_4(\gamma))=\{0,1,2,\infty\}$ and*

$$M_4(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f})(\infty)=\big(S^2,(a,b),(d,-c),(e,f)\big), \tag{9}$$

$$M_4(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f})(2)=\big(D,(a-b,b),(e-f,f)\big)\bigcup_{\begin{pmatrix}0&1\\1&0\end{pmatrix}}\big(D,(c,d),(2,-1)\big), \tag{10}$$

$$M_4(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f})(1)=\big(S^2,(a-2b,b),(c-d,c),(e-2f,f)\big), \tag{11}$$

$$M_4(\tfrac{a}{b},\tfrac{c}{d},\tfrac{e}{f})(0)=\big(D,(f,-e),(b,2b-a)\big)\bigcup_{\begin{pmatrix}0&1\\1&0\end{pmatrix}}\big(D,(2,1),(c-2d,d)\big), \tag{12}$$

or $5\leqslant e_\tau(M_4(\gamma))\leqslant 6$ and

- *$M_4(\gamma)\cong M_4(\alpha)$ for some α in Tables* 5-6*, in Section 4, and with respect to the basis induced from M_4 we have*

$$E_\tau(M_4(\alpha))=\begin{cases}\{\beta_1,\beta_2,0,1,2,\infty\} & \textit{if } \alpha \textit{ is found in Table } 5;\\ \{\beta,0,1,2,\infty\} & \textit{otherwise,}\end{cases}$$

  where β,β_1,β_2 are found in Tables 5 *and* 6*.*
- *Identities* (9)-(12) *hold, and the $M_5(\alpha)(\beta_i)$, $M_4(\alpha)(\beta)$ are shown in Tables* 5 *and* 6*.*

While proving Theorems 3 and 4 we will often see identities between fillings of the M_i. When a homeomorphism $M_5(\alpha)\simeq M_5(\beta)$ follows as a consequence of maps (1)-(5) we will write $M_5(\alpha)\sim M_5(\beta)$. In addition to realising homeomorphisms between fillings of M_5, we will often use the Rolfsen twist (see [Rol]) to obtain homeomorphisms between fillings of the M_i; when $M_5(\alpha)\simeq M_k(\beta)$ is induced from a (composition of) Rolfsen twist(s), we will write $M_5(\alpha)\simeq M_k(\beta)$.

Proof of Theorem 3. By (1)-(5), we have

$$[\![(1)]\!]=[(1)]\sqcup[(\infty)]\sqcup[(0)]; \tag{13}$$

$$[\![(-1)]\!] = [(-1)] \sqcup [(\tfrac{1}{2})] \sqcup [(2)]; \tag{14}$$

$$[\![(-2)]\!] = [(-2)] \sqcup [(-\tfrac{1}{2})] \sqcup [(\tfrac{1}{3})] \sqcup [(\tfrac{2}{3})] \sqcup [(\tfrac{3}{2})] \sqcup [(3)]. \tag{15}$$

Theorem 2 implies that all slopes in $[\![(1)]\!]$ are exceptional, moreover, any instruction containing a slope in $[\![(-1)]\!]$ factors through M_4. So for τ a boundary component of a hyperbolic $M_5(\gamma)$ we have $\{0, 1, \infty\} \subseteq E_\tau(M_5(\gamma))$ and $e_\tau(M_5(\gamma)) \geq 3$; when γ does not factor through M_4 no slope in $[\![(-1)]\!]$ is contained in γ. Identity (6) is established in [MPR], and Identities (7) and (8) are established using (1)-(3) and (6). We will now describe up to (1)-(3) all hyperbolic γ's not factoring through M_4 with $e_\tau(M_5(\gamma)) > 3$. For such γ, using (1)-(2), we may assume that τ corresponds to the neighbourhood of the fourth chain component of 5CL.

So we suppose that $\gamma_0, \gamma_1, \gamma_2, \gamma_3 \in \mathbb{Q} \cup \{\varnothing\}$, with $\gamma = (\gamma_0, \gamma_1, \gamma_2, \gamma_3)$ a hyperbolic filling instruction on M_5 that does not factor through M_4. If β is an exceptional slope on $M_5(\gamma)$ then, up to (1)-(3), $(\gamma_0, \gamma_1, \gamma_2, \gamma_3, \beta)$ contains one of the seven isolated exceptional filling instruction from Theorem 2. If $\beta \neq 0, 1, \infty$ then any such isolated exceptional filling instruction contains at most one slope in $[\![(-1)]\!]$, and contains two slopes in $[\![(-2)]\!]$ (see Theorem 2 and (15)). Thus at least one of the slopes in γ belongs to $[\![(-2)]\!]$. It is a routine consequence of maps (1)-(3) that we may assume without loss of generality that $\alpha_0 = -2$ and that τ remains the boundary corresponding to the neighbourhood of the 4th chain component of 5CL.

So for γ hyperbolic, not factoring through M_4, we have $[\![\gamma]\!] = [\![\alpha]\!]$ where $\alpha_i \notin \{-1, \frac{1}{2}, 2\}$ and $\alpha_0 = -2$. We define (α, β) to be $(\alpha_0, \alpha_1, \alpha_2, \alpha_3, \beta)$. If β is an exceptional slope of $M_5(\alpha)$ then (α, β) contains an isolated exceptional filling instruction by Theorem 2. Moreover, using Theorem 2, if (α, β) contains an isolated exceptional filling instruction then either β is a slope in $[\![(1)]\!]$, or β is a slope in $[\![(-1)]\!]$ and (α, β) factors through M_3, or $M_5(\alpha, \beta) \in \{m_1, m_2, m_3\}$. As we are assuming that α does not factor through M_4, if (α, β) factors through M_3 then we require $\beta \in \{-1, \frac{1}{2}, 2\}$.

We define the following sets of (α, β)'s;

- We define l to be the set of exceptional (α, β)'s such that $M_5(\alpha, \beta)$ is in $\{m_1, m_2, m_3\}$;
- We define l_{-1} to be the set of exceptional (α, β)'s factoring through M_3 with $\beta = -1$;

- We define $l_{\frac{1}{2}}$ to be the set of exceptional (α, β)'s factoring through M_3 with $\beta = \frac{1}{2}$;
- We define l_2 to be the set of exceptional (α, β)'s factoring through M_3 with $\beta = 2$;

We know that, every γ with $M_5(\gamma)$ a hyperbolic manifold, γ not factoring through M_4, and $e_\tau(M_5(\gamma)) > 3$ has $[\![\gamma]\!] = [\![\alpha]\!] = [\![(\alpha_0, \alpha_1, \alpha_2, \alpha_3)]\!]$ where $\alpha_0 = -2$. The set $l_{-1} \cup l_{\frac{1}{2}} \cup l_2$ is the set of all exceptional filling instructions (α, β) factoring through M_3 with $M_5(\alpha)$ hyperbolic not factoring through M_4 and $\alpha_0 = -2$, and the set l is the set of all exceptional filling instructions (α, β) not factoring through M_3 with $\alpha_0 = -2$.

So, up to (1)-(3) every hyperbolic filling instruction γ not factoring through M_4 with $e_\tau(M_5(\gamma)) > 3$ is equivalent to a filling instruction α of the form $(-2, \alpha_1, \alpha_2, \alpha_3)$, every $(-2, \alpha_1, \alpha_2, \alpha_3)$ with $e_\tau(M_5(-2, \alpha_1, \alpha_2, \alpha_3) > 3$ and $(-2, \alpha_1, \alpha_2, \alpha_3)$ not factoring through M_4 is some α in $l \cup l_{-1} \cup l_{\frac{1}{2}} \cup l_2$, and every exceptional slope on $M_5(-2, \alpha_1, \alpha_2, \alpha_3)$ not in $\{0, 1, \infty\}$ is some β in $l \cup l_{-1} \cup l_{\frac{1}{2}} \cup l_2$. Now we can easily describe, up to $\sim$, all such α with more than 3 exceptional slopes.

Let $p : l \cup l_{-1} \cup l_{\frac{1}{2}} \cup l_2 \to A$ be defined by $(\alpha, \beta) \mapsto \alpha$. For α in $p(l \cup l_{-1} \cup l_{\frac{1}{2}} \cup l_2)$ define B_α to be the set of all β's such that (α, β) is contained in $l \cup l_{-1} \cup l_{\frac{1}{2}} \cup l_2$. It is clear that $E_\tau\big(M_5(\alpha)\big) = \{0, 1, \infty\} \cup B_\alpha$ and that $\{\big(M_5(\alpha), E_\tau(M_5(\alpha))\big)\}_\alpha$ is a complete list of all $\big(M_5(\alpha), E_\tau(M_5(\alpha))\big)$ pairs with $\alpha = (-2, \alpha_1, \alpha_2, \alpha_3)$, $M_5(\alpha)$ hyperbolic and $e_\tau\big(M_5(\alpha)\big) > 3$. We now explicitly construct the sets $l, l_{-1}, l_{\frac{1}{2}}, l_2$.

Construction of the set l: We can see using maps (1)-(5) that every (α, β) with $M_5(\alpha, \beta) = m_i$ with $\alpha_0 = -2$ and α hyperbolic not factoring through M_4 is contained in the following set

$$l = \Big\{\big(-2, \tfrac{1}{4}, \tfrac{3}{2}, \tfrac{4}{3}, \tfrac{1}{2}\big), \big(-2, -\tfrac{1}{2}, 3, 3, -\tfrac{1}{2}\big), \big(-2, \tfrac{1}{3}, 3, \tfrac{1}{3}, -2\big), \big(-2, -2, \tfrac{1}{3}, 3, \tfrac{1}{3}\big),$$
$$\big(-2, -\tfrac{1}{2}, -2, \tfrac{3}{2}, \tfrac{3}{2}\big), \big(-2, \tfrac{3}{2}, \tfrac{3}{2}, -2, -\tfrac{1}{2}\big), \big(-2, -2, -2, -2, -2\big), \big(-2, \tfrac{1}{3}, \tfrac{3}{2}, \tfrac{3}{2}, \tfrac{1}{3}\big)\Big\}.$$

Before constructing the sets $l_{-1}, l_{\frac{1}{2}}, l_2$ we make some initial remarks. It is easy to see that if a surgery instruction on 5CL contains $((-1)_0, (-1)_2)$, then it factors through M_3. Up to the action of maps (1)-(5) the surgery instruction $((-1)_0, (-1)_2)$ is equivalent to the following surgery instructions:

$$\Big\{[((-1)_0, (-2)_1)], [((\tfrac{1}{2})_0, (\tfrac{2}{3})_2)], [((\tfrac{1}{2})_0, 3_1)], [((\tfrac{1}{2})_0, (\tfrac{3}{2})_1)], [((\tfrac{2}{3})_0, 2_1)],$$
$$[((\tfrac{1}{2})_0, (\tfrac{1}{3})_2)], [(2_0, (-\tfrac{1}{2})_2)], [((\tfrac{1}{3})_0, 2_1)], [((-1)_0, 3_2)], [((-1)_0, (\tfrac{3}{2})_2)],$$
$$[(2_0, (-2)_2)], [((-1)_0, (-\tfrac{1}{2})_1)], [((-1)_0, (-1)_2)], [((-1)_0, (\tfrac{1}{2})_2)],$$

$$[((\tfrac{1}{2})_0,(2)_2)],[((-1)_0,(2)_1)],[((-1)_0,(\tfrac{1}{2})_1)],[((2)_0,(2)_2)]\Big\}$$

We will now see that the requirements of α being hyperbolic and not factoring through M_4 allow us to completely construct $l_{-1}, l_{\frac{1}{2}}, l_2$.

Construction of the set l_{-1}: In this case, for $\alpha = (-2, \alpha_1, \alpha_2, \alpha_3)$, we have $M_5(\alpha)(-1) \simeq M_3(\alpha_1 + 1, \alpha_2, \alpha_3 + 2)$, and (α, β) is an exceptional filling of M_5 if and only if $(\alpha_1 + 1, \alpha_2, \alpha_3 + 2)$ contains an isolated exceptional filling instruction on M_3; i.e. using the classification in [MP] we can see that -1 is an exceptional slope on $M_5(\alpha)$ if $(\alpha_1 + 1, \alpha_2, \alpha_3 + 2)$ contains an isolated instruction in

$$\Big\{0,1,2,3,\infty,(-1,-1),(4,\tfrac{1}{2}),(\tfrac{3}{2},\tfrac{5}{2}),(5,5,\tfrac{1}{2}),(4,4,\tfrac{2}{3}),(4,\tfrac{3}{2},\tfrac{3}{2}),(4,\tfrac{1}{3},-1),(\tfrac{8}{3},\tfrac{3}{2},\tfrac{3}{2}),$$
$$(\tfrac{5}{2},\tfrac{5}{2},\tfrac{4}{3}),(\tfrac{5}{2},\tfrac{5}{3},\tfrac{5}{3}),(\tfrac{7}{3},\tfrac{7}{3},\tfrac{3}{2}),(-1,-2,-2),(-1,-2,-3),(-1,-2,-4),(-1,-2,-5),$$
$$(-1,-3,-3),(-2,-2,-2)\Big\},$$

For α non-exceptional and not factoring through M_4, no slope in α is in $[(1)]$ or $[(-1)]$; that is $\alpha_1 + 1 \notin \{\infty, 2, 1, 0, \frac{3}{2}, 3\}$, $\alpha_2 \notin \{\infty, 1, 0, -1, \frac{1}{2}, 2\}$, $\alpha_3 + 2 \notin \{\infty, 3, 2, 1, \frac{5}{2}, 4\}$. With these conditions it is easy to see that β is an exceptional slope on a hyperbolic $M_5(\alpha)$ if and only if one of the following holds:

- $\alpha_2 = 3$
- $\alpha_3 + 2 = 0$
- $(\alpha_1 + 1, \alpha_3 + 2)$ belongs to $\{(-1,-1),(\frac{5}{2},\frac{3}{2}),(4,\frac{1}{2})\}$
- $(\alpha_2, \alpha_1 + 1)$ belongs to $\{(\frac{3}{2},\frac{5}{2}),(4,\frac{1}{2})\}$
- $(\alpha_2, \alpha_3 + 2)$ belongs to $\{(\frac{5}{2},\frac{3}{2}),(4,\frac{1}{2})\}$
- $(\alpha_1 + 1, \alpha_2, \alpha_3 + 2)$ belongs to

$$\Big\{(5,5,\tfrac{1}{2}),(\tfrac{1}{2},5,5),(4,4,\tfrac{2}{3}),(4,\tfrac{3}{2},\tfrac{3}{2}),(4,\tfrac{1}{3},-1),(\tfrac{1}{3},4,-1),(-1,4,\tfrac{1}{3}),(\tfrac{8}{3},\tfrac{3}{2},\tfrac{3}{2}),$$
$$(\tfrac{5}{2},\tfrac{5}{2},\tfrac{4}{3}),(\tfrac{5}{2},\tfrac{5}{3},\tfrac{5}{3}),(\tfrac{5}{3},\tfrac{5}{2},\tfrac{5}{3}),(\tfrac{7}{3},\tfrac{7}{3},\tfrac{3}{2}),(\tfrac{7}{3},\tfrac{3}{2},\tfrac{7}{3}),(-1,-2,-2),(-2,-2,-1),$$
$$(-1,-2,-3),(-3,-2,-1),(-2,-3,-1),(-1,-3,-2),(-1,-2,-4),$$
$$(-4,-2,-1),(-1,-4,-2),(-2,-4,-1),(-1,-2,-5),(-5,-2,-1),$$
$$(-1,-5,-2),(-2,-5,-1),(-1,-3,-3),(-3,-3,-1),(-2,-2,-2))\Big\}.$$

Thus, the set of all $(-2, \alpha_1, \alpha_2, \alpha_3, -1)$ constructed in the above analysis is the set l_{-1}.

Construction of the set $l_{\frac{1}{2}}$: Reasoning as in the case $\beta = -1$ we see (α, β) factors through M_3 when one of α_1, α_2 is in $\{\frac{1}{3}, \frac{2}{3}\}$ or $\alpha_3 \in \{3, \frac{3}{2}\}$ (see $[\![(-1_0, -1_2)]\!]$ shown above). We examine each of these 6 cases individually and enumerate all $(-2, \alpha_1, \alpha_2, \alpha_3, \frac{1}{2})$ with the desired properties.

If $\alpha_3 = 3$ then

$$M_5\big(-2, \alpha_1, \alpha_2, 3, \tfrac{1}{2}\big) \sim M_5\big(-1, 3, \alpha_2^{-1}, \alpha_1^{-1}, -2\big) \simeq M_3\big(5, \alpha_2^{-1}, \alpha_1^{-1} + 1\big).$$

The result is that $\frac{1}{2}$ is an exceptional slope on $M_5\big(-2, \alpha_1, \alpha_2, \alpha_3\big)$ if and only if one of the following holds; $\alpha_1^{-1} = 3$, $\big(\alpha_2^{-1}, \alpha_1^{-1} + 1\big) = \big(\frac{3}{2}, \frac{5}{2}\big)$, $(4, \frac{1}{2})$, $(5, \frac{1}{2})$.

If $\alpha_3 = \frac{3}{2}$ then

$$\begin{aligned} M_5\big(-2, \alpha_1, \alpha_2, \tfrac{3}{2}, \tfrac{1}{2}\big) &\sim M_5\big(-2, -2, (1 - \alpha_2)^{-1}, 1 - \alpha_1^{-1}, -1\big) \\ &\simeq M_3\big(-1, (1 - \alpha_2)^{-1}, 3 - \alpha_1^{-1}\big). \end{aligned}$$

We conclude that $\frac{1}{2}$ is an exceptional slope on $M_5\big(-2, \alpha_1, \alpha_2, \alpha_3\big)$ if and only if one of the following holds; $(1-\alpha_2)^{-1} = 3$, $3-\alpha_1^{-1} = 0, -1$, $((1-\alpha_2)^{-1}, 3-\alpha_1^{-1}\big) = (\frac{5}{2}, \frac{3}{2})$, $(4, \frac{1}{2})$, $(-2, -2)$, $(-2, -3)$, $(-3, -2)$, $(-2, -4)$, $(-2, -5)$, $(-5, -2)$, $(4, \frac{1}{3})$.

If $\alpha_1 = \frac{1}{3}$ then

$$\begin{aligned} M_5(\alpha)(\beta) &\sim M_5\big((1 - \alpha_3)^{-1}, -2, (1 - \alpha_2)^{-1}, -2, -1\big) \\ &\simeq M_3\big(2 + (1 - \alpha_3)^{-1}, -2, 1 + (1 - \alpha_2)^{-1}\big). \end{aligned}$$

The result is that $\frac{1}{2}$ is an exceptional slope on $M_5\big(-2, \alpha_1, \alpha_2, \alpha_3\big)$ if and only if one of the following holds; $1 + (1 - \alpha_2)^{-1} = 3$, $2 + (1 - \alpha_3)^{-1} = 0$, $(2 + (1 - \alpha_3)^{-1}, 1 + (1 - \alpha_2)^{-1}) = (-1, -1)$, $(\frac{1}{2}, 4)$, $(\frac{3}{2}, \frac{5}{2})$.

If $\alpha_1 = \frac{1}{3}$ then

$$M_5\big(-2, \alpha_1, \alpha_2, \alpha_3, \tfrac{1}{2}\big) \sim M_5\big(\tfrac{1}{3}, -1, -2, (1-\alpha_1)^{-1}, \alpha_3\big) \simeq M_3\big(\tfrac{7}{3}, 1+(1-\alpha_1)^{-1}, \alpha_3\big).$$

We see that $\frac{1}{2}$ is an exceptional slope on $M_5\big(-2, \alpha_1, \alpha_2, \alpha_3\big)$ if and only if one of the following holds; $\alpha_3 = 3$, $(\alpha_3, 1 + (1 - \alpha_1)^{-1})$=$(\frac{3}{2}, \frac{5}{2})$, $(\frac{3}{2}, \frac{7}{3})$.

If $\alpha_1 = \frac{2}{3}$ then

$$\begin{aligned} M_5(\alpha)(\beta) &\sim M_5\big(\tfrac{2}{3}, (1 - \alpha_2^{-1})^{-1}, -1, -2, (1 - \alpha_3^{-1})^{-1}\big) \\ &\simeq M_3\big(\tfrac{2}{3}, 2 + (1 - \alpha_2^{-1})^{-1}, 1 + (1 - \alpha_3^{-1})^{-1}\big). \end{aligned}$$

We find that $\frac{1}{2}$ is an exceptional slope on $M_5\big(-2,\alpha_1,\alpha_2,\alpha_3\big)$ if and only if one of the following holds; $2+(1-\alpha_2^{-1})^{-1}=0$, $(2+(1-\alpha_2^{-1})^{-1},1+(1-\alpha_3^{-1})^{-1})=(-1,-1)$, $(\frac{3}{2},\frac{5}{2})$.

If $\alpha_2=\frac{2}{3}$ then

$$\begin{aligned} M_5(\alpha)(\beta) &\sim M_5\big(\tfrac{2}{3},-2,-1,(1-\alpha_1^{-1})^{-1},(1-\alpha_3^{-1})^{-1}\big) \\ &\simeq M_3\big(\tfrac{5}{3},2+(1-\alpha_1^{-1})^{-1},(1-\alpha_3^{-1})^{-1}\big). \end{aligned}$$

The result is that $\frac{1}{2}$ is an exceptional slope on $M_5\big(-2,\alpha_1,\alpha_2,\alpha_3\big)$ if and only if one of the following holds; $(1-\alpha_3^{-1})^{-1}=3$, $(2+(1-\alpha_1^{-1})^{-1},(1-\alpha_3^{-1})^{-1})=(\frac{3}{2},\frac{5}{2})$, $(\frac{1}{2},4)$, $(\frac{5}{3},\frac{5}{2})$.

Thus, the set of all $(-2,\alpha_1,\alpha_2,\alpha_3,\frac{1}{2})$ constructed in the above analysis is the set $l_{\frac{1}{2}}$.

Construction of the set l_2: For $(\alpha,\beta)=(-2,\alpha_1,\alpha_2,\alpha_3,2)$ to factor through M_3 we need one of α_1,α_2 to be in $\{-2,-\frac{1}{2}\}$ or $\alpha_3\in\{\frac{1}{3},\frac{2}{3}\}$ (see $\llbracket(-1_0,-1_2)\rrbracket$ shown above). We construct all (α,β) for each of these 6 cases individually.

If $\alpha_1=-2$ then

$$M_5(\alpha)(2)\sim M_5\big(\tfrac{3}{2},1-\alpha_2^{-1},(1-\alpha_3)^{-1},-2,-1\big)\simeq M_3\big(\tfrac{7}{2},1-\alpha_2^{-1},1+(1-\alpha_3)^{-1}\big).$$

When no α_1, α_2, α_3 is a slope in $\llbracket(1)\rrbracket$, $\llbracket(-1)\rrbracket$ we have $1-\alpha_2^{-1}\notin\{\infty,0,1,2,\frac{1}{2},-1\}$ and $1+(1-\alpha_3)^{-1}\notin\{2,\infty,1,\frac{3}{2},0,3\}$. From [MP] we see that 2 is exceptional if and only if one of the following holds; $1-\alpha_2^{-1}=3$, $(1-\alpha_2^{-1},1+(1-\alpha_3)^{-1})=(\frac{3}{2},\frac{5}{2})$, $(4,\frac{1}{2})$.

If $\alpha_2=-2$ then

$$M_5(\alpha)(\beta)\sim M_5\big(-1,-2,\tfrac{1}{3},1-\alpha_1^{-1},1-\alpha_3^{-1}\big)\simeq M_3\big(\tfrac{4}{3},1-\alpha_1^{-1},3-\alpha_1^{-1}\big).$$

Using [MP] we get that 2 is exceptional on $M_5(\alpha)$ if and only if one of the following holds; $1-\alpha_1^{-1}=3$, $3-\alpha_3^{-1}=0$, $(1-\alpha_1^{-1},3-\alpha_3^{-1})=(4,\frac{1}{2})$, $(\frac{5}{2},\frac{3}{2})$.

If $\alpha_1=-\frac{1}{2}$ then

$$M_5(\alpha)(\beta)\sim M_5\big((\alpha_3-1)^{-1},1-\alpha_1,-\tfrac{1}{2},-2,-1\big)\simeq M_3\big(2+(\alpha_3-1)^{-1},1-\alpha_1,\tfrac{1}{2}\big).$$

We find that 2 is exceptional on $M_5(\alpha)$ if and only if one of the following holds; $1-\alpha_1=3$, $(1-\alpha_1,2+(\alpha_3-1)^{-1})=(4,\frac{1}{2})$, $(\frac{5}{2},\frac{3}{2})$.

If $\alpha_2=-\frac{1}{2}$ then

$$M_5(\alpha)(\beta)\sim M_5\big(-2,-1,\tfrac{2}{3},1-\alpha_1,\alpha_3^{-1}\big)\simeq M_3\big(\tfrac{8}{3},1-\alpha_1,1+\alpha_3^{-1}\big).$$

We observe that 2 is exceptional on $M_5(\alpha)$ if and only if one of the following holds; $1-\alpha_1 = 3$, $(1-\alpha_1, 1+\alpha_3^{-1}) = (4, \frac{1}{2})$, $(\frac{3}{2}, \frac{5}{2})$.

If $\alpha_3 = \frac{2}{3}$ then

$$M_5(\alpha)(\beta) \sim M_5\big(-2, 1-\alpha_2, -\tfrac{1}{2}, \alpha_1^{-1}, -1\big) \simeq M_3\big(2-\alpha_2, -\tfrac{1}{2}, 2+\alpha_1^{-1}\big).$$

We get that 2 is exceptional on $M_5(\alpha)$ if and only if one of the following holds; $2+\alpha_1^{-1} = 0$, $(1-\alpha_2, \alpha_1^{-1}) = (-1,-1)$, $(2-\alpha_2, 2+\alpha_1^{-1}) = (4, \frac{1}{2})$, $(\frac{5}{2}, \frac{3}{2})$.

If $\alpha_3 = \frac{1}{3}$ then

$$M_5(\alpha)(\beta) \sim M_5\big(-1, \alpha_2, \tfrac{1}{3}, 1-\alpha_1^{-1}, -2\big) \simeq M_3\big(\alpha_2+2, \tfrac{1}{3}, 2-\alpha_1^{-1}\big).$$

The result is that 2 is an exceptional slope on $M_5(\alpha)$ if and only if one of the following holds; $\alpha_2 + 2 = 0$, $(\alpha_2+2, 2-\alpha^{-1}) = (\frac{1}{2}, 4)$, $(\frac{3}{2}, \frac{5}{2})$, $(-1,-1)$, $(-1,4)$. Thus, the set of all $(-2, \alpha_1, \alpha_2, \alpha_3, 2)$ constructed in the above analysis is the set l_2.

This completes the construction of $l \cup l_{-1} \cup l_{\frac{1}{2}} \cup l_2$ and the sets $\{(M_5(\alpha), E_\tau(M_5(\alpha))\}$. The last step is to reduce the size of $\{(M_5(\alpha), E_\tau(M_5(\alpha))\}$ using maps (1)-(5). The result is shown in Tables 1 and 4. The corresponding exceptional manifolds shown in Tables 1 and 4 are written down using the description of isolated exceptional fillings of M_3 and M_5 provided in [MP] and [MPR] respectively. □

The proof of Theorem 4 follows a similar line of reasoning:

Proof of Theorem 4. We let $\gamma = (\gamma_0, \gamma_1, \gamma_2, \gamma_3)$ be a hyperbolic filling instruction on M_4 with at least on $\varnothing$ slope. We translate the problem on 4CL into a problem on 5CL; using the Rolfsen twist we see that $M_4(\gamma) \simeq M_5(\delta)$ where $\delta = (\gamma_0 - 1, \gamma_1, \gamma_2, \gamma_3 - 1, -1)$ (here $\gamma_0 - 1$, $\gamma_3 - 1$ should be read as $\varnothing$ when γ_0, $\gamma_3 = \varnothing$). Now the argument proceeds exactly as in the proof of Theorem 3.

We use Identities (1)-(3) to assume without loss of generality that $\delta_0 = -1$ and that τ is the boundary corresponding to the neighbourhood of the 4th chain component of 5CL; namely we can assume $\delta = (\delta_0, \delta_1, \delta_2, \delta_3, \varnothing)$ is a hyperbolic filling instruction on M_5 that does not factor through M_3, and that some slope of δ is in $\{-1, \frac{1}{2}, 2\}$.

The conditions of δ being non-exceptional and not factoring through M_3 means that $\delta_1 \notin \{0, 1, \infty, -1, -2, -\frac{1}{2}\}$ and $\delta_2, \delta_3 \notin \{0, 1, \infty, -1, \frac{3}{2}, 3\}$. Theorem 2 implies that every $\beta \in \{-1, 0, 1, \infty\}$ is an exceptional slope on $M_5(\delta)$, and so $\{-1, 0, 1, \infty\} \subseteq E_\tau(M_5(\delta))$ and $e_\tau(M_5(\delta)) \geq 4$. Identities (9)-(11) follow directly from Identities (6)-(8) while (12) is established using (4) and then a composition of (1)-(3). More generally, Theorem 2 implies that if β is an exceptional slope on $M_5(\delta)$ then $\beta \in \{-1, 0, 1, \infty\}$, or (δ, β) factors through M_3, or $M_5(\delta, \beta) = m_2$. The (δ, β) coming from m_2 are given by $(-1, -3, -2, -2, -3)$, $(-1, \frac{1}{3}, \frac{4}{3}, \frac{4}{3}, \frac{1}{3})$, $(-1, -\frac{1}{3}, 4, \frac{2}{3}, 3)$, $(-1, 3, \frac{2}{3}, 4, -\frac{1}{3})$. This implies that β is in one of $[\![(1)]\!]$, $[\![(-1)]\!]$, $[\![(-2)]\!]$, $[\![(-3)]\!]$.

Every slope in $\{[\![(-1)]\!], [\![(-2)]\!], [\![(-3)]\!]\}$ is examined individually as in the proof of Theorem 3 to obtain a complete list of all $\big(M_5(\delta), E_\tau(M_5(\delta))\big)$ pairs that have $\delta = \big(-1, \frac{p}{q}, \frac{r}{s}, \frac{u}{v}, \varnothing\big)$ hyperbolic not factoring through M_3 with $e\big(M_5(\delta)\big) > 4$.

Lastly, the list of δ produced is reduced using the maps (1)-(5), and a positive twist about the zeroth component of the 5CL gives the list of α's on M_4 and the exceptional slopes β_i on $M_4(\alpha)$ shown in Table 5-6. The corresponding exceptional manifolds shown in Table 5-6 are written down using the description of exceptional fillings on M_3 and M_5 provided in [MP] and [MPR]. $\square$

We finish by pointing out that Theorems 3 and 4 imply that any one-cusped hyperbolic manifold M obtained by surgery on 5CL with 10 exceptional slopes, or having two exceptional slopes at distance 8, is obtained by surgery on 3CL and that the only such M's obtained by surgery on 3CL are the figure-8 knot exterior and its sister (see the Appendix in [MP]) which shows the final two assertions of Theorem 1. The second assertion in Theorem 1 is a simple consequence of the fact that, up to (1)-(5), the only $M_5(\alpha)$ with a pair of exceptional slopes at distance 4 with α not factoring through M_3 is $M_4(-2, -\frac{1}{2}, -2)$ with the slopes $-1, 3$ (see Tables 2-1), and $M_4(-2, -\frac{1}{2}, -2)(-1)$, $M_4(-2, -\frac{1}{2}, -2)(3)$ are not lens spaces; all $M_3(\alpha)$ with a pair of exceptional slopes at distance 4 can be found in [MP], and it is not hard to check that the corresponding pair of exceptional fillings are never a lens space, toroidal pair. Lastly, we remark that the description of exceptional fillings provided in Theorems 3 and 4, and [MP] can be used to easily enumerate all examples of exceptional pairs at maximal distance; one such enumeration shows that no counterexample to the Cabling conjecture is obtained by surgery on 5CL (see [Rou] for full details).

4. TABLES

α	Additional exceptional slopes β_i	Exceptional filling $M_5(\alpha)(\beta_i)$
$(-2,-\frac{1}{2},3,3)$	$\beta_1=-1$ $\beta_2=-\frac{1}{2}$	$\big(S^2,(2,1),(5,-2),(4,-1)\big)$ $\big(A,(2,-1)\big)\big/_{\begin{pmatrix}1&2\\1&1\end{pmatrix}}$
$(-2,\frac{3}{2},\frac{3}{2},-2)$	$\beta_1=-1$ $\beta_2=-\frac{1}{2}$	$\big(\mathbb{RP}^2,(2,-1),(3,1)\big)$ $\big(A,(2,-1)\big)\big/_{\begin{pmatrix}1&2\\1&1\end{pmatrix}}$
$(-2,-3,-\frac{1}{2},-2)$	$\beta_1=-1$ $\beta_2=2$	$\big(S^2,(2,-1),(3,1),(9,2)\big)$ $\big(D,(2,1),(3,1)\big)\bigcup_{\begin{pmatrix}1&1\\1&0\end{pmatrix}}\big(D,(2,1),(3,-8)\big)$
$(-2,-\frac{1}{3},3,\frac{2}{3})$	$\beta_1=-1$ $\beta_2=2$	$\big(S^2,(2,1),(7,-2),(5,-3)\big)$ $\big(A,(2,3)\big)\big/_{\begin{pmatrix}0&1\\1&0\end{pmatrix}}$
$(-2,-\frac{1}{2},3,\frac{2}{3})$	$\beta_1=-1$ $\beta_2=2$	$\big(S^2,(2,1),(5,-2),(5,-3)\big)$ $\big(S^2,(2,-1),(3,1),(11,2)\big)$
$(-2,-2,-2,-2)$	$\beta_1=-1$ $\beta_2=-2$	$\big(S^2,(2,-1),(3,1),(7,1)\big)$ $\big(D,(2,1),(2,-1)\big)\bigcup_{\begin{pmatrix}-1&5\\1&-4\end{pmatrix}}\big(D,(2,1),(3,1)\big)$

TABLE 1. α a filling instruction on M_5 not factoring through M_4 with $e_\tau\big(M_5(\alpha)\big)=5$, and $E_\tau\big(M_5(\alpha)\big)=\{\beta_1,\beta_2,0,1,\infty\}$.

α	Additional exceptional slopes β	Exceptional filling $M_5(\alpha)(\beta)$
$(-2, \frac{p}{q}, 3, \frac{u}{v})$	-1	$\bigl(D, (2,1), (p,-q)\bigr) \bigcup_{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}} \bigl(D, (2,1), (u+v,-v)\bigr)$
$(-2, \frac{p}{q}, \frac{r}{s}, -2)$	-1	$\bigl(D, (s,2s-r), (q,q-p)\bigr) \bigcup_{\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}} \bigl(D, (2,1), (3,1)\bigr)$
$(-2, \frac{3}{2}, \frac{3}{2}, \frac{u}{v})$	-1	$\bigl(D, (2,1), (3,1)\bigr) \bigcup_{\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}} \bigl(D, (2,1), (u,-v)\bigr)$
$(-2, \frac{p}{q}, \frac{5}{2}, -\frac{1}{2})$	-1	$\bigl(D, (2,1), (3,1)\bigr) \bigcup_{\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}} \bigl(D, (2,1), (q-p,q)\bigr)$
$(-2, -2, \frac{r}{s}, -3)$	-1	$\bigl(A, (s,s-r)\bigr) /_{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}$
$(-2, -\frac{1}{2}, 4, \frac{u}{v})$	-1	$\bigl(D, (2,1), (3,1)\bigr) \bigcup_{\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}} \bigl(D, (2,1), (v,-u-2v)\bigr)$
$(-2, \frac{p}{q}, 4, -\frac{3}{2})$	-1	$\bigl(D, (2,1), (3,1)\bigr) \bigcup_{\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}} \bigl(D, (2,1), (q,-p-q)\bigr)$
$(-2, \frac{1}{3}, \frac{r}{s}, \frac{3}{2})$	$\frac{1}{2}$	$\bigl(S^2, (2,-1), (3,1), (4s+r,s)\bigr)$

TABLE 2. α a filling instruction on M_5 not factoring through M_4 with $e_\tau\bigl(M_5(\alpha)\bigr) = 4$, and $E_\tau\bigl(M_5(\alpha)\bigr) = \{\beta, 0, 1, \infty\}$, part 1/3.

α	Additional exceptional slopes β	Exceptional filling $M_5(\alpha)(\beta)$
$(-2,4,5,-\frac{3}{2})$ $(-2,-\frac{1}{2},5,3)$	-1	$\big(A,(2,1)\big)/\begin{pmatrix}0&1\\1&0\end{pmatrix}$
$(-2,3,4,-\frac{4}{3})$	-1	$\big(A,(2,1)\big)/\begin{pmatrix}1&1\\1&0\end{pmatrix}$
$(-2,3,\frac{3}{2},-\frac{1}{2})$	-1	$T/\begin{pmatrix}-3&1\\-1&0\end{pmatrix}$
$(-2,-2,4,-\frac{5}{3})$ $(-2,-\frac{2}{3},4,-3)$	-1	$\big(D,(2,1),(2,1)\big)\bigcup_{\begin{pmatrix}0&1\\-1&-1\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
$(-2,\frac{2}{3},\frac{5}{2},-\frac{1}{3})$	-1	$\big(D,(2,1),(2,1)\big)\bigcup_{\begin{pmatrix}-1&1\\0&-1\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
$(-2,\frac{4}{3},\frac{3}{2},\frac{1}{3})$	-1	$\big(A,(2,1)\big)/\begin{pmatrix}1&1\\1&0\end{pmatrix}$
$(-2,-3,-2,-3)$ $(-2,-2,-2,-4)$ $(-2,-2,-2,-4)$	-1	$\big(S^2,(2,-1),(3,1),(7,1)\big)$

TABLE 3. α a filling instruction on M_5 not factoring through M_4 with $e\big(M_5(\alpha)\big)=4$, and $E_\tau\big(M_5(\alpha)\big)=\{\beta,0,1,\infty\}$, part 2/3.

α	Additional exceptional slope β	Exceptional filling $M_5(\alpha)(\beta)$
$(-2,-4,-2,-3)$ $(-2,-2,-2,-5)$ $(-2,-3,-3,-3)$ $(-2,-2,-3,-4)$	-1	$\big(S^2,(2,-1),(3,1),(5,1)\big)$
$(-2,-4,-3,-3)$ $(-2,-2,-3,-5)$ $(-2,-2,-3,-5)$ $(-2,-4,-3,-3)$	-1	$\big(D,(2,1),(2,1)\big)\bigcup_{\begin{pmatrix}1&2\\0&-1\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
$(-2,-2,-4,-4)$ $(-2,-5,-2,-3)$ $(-2,-3,-4,-3)$ $(-2,-2,-2,-6)$	-1	$\big(S^2,(2,-1),(4,1),(5,1)\big)$
$(-2,-2,-2,-7)$ $(-2,-6,-2,-3)$ $(-2,-3,-5,-3)$ $(-2,-2,-5,-4)$	-1	$\big(D,(2,1),(2,1)\big)\bigcup_{\begin{pmatrix}0&1\\1&0\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
$(-2,-3,-2,-4)$	-1	$\big(D,(2,1),(2,1)\big)\bigcup_{\begin{pmatrix}2&3\\-1&-2\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
$(-2,\frac{3}{2},\frac{5}{2},-\frac{2}{3})$	-1	$\big(A,(2,1)\big)/_{\begin{pmatrix}2&1\\1&0\end{pmatrix}}$
$(-2,-2,\frac{1}{4},3)$	$\frac{1}{2}$	$\big(S^2,(2,1),(3,-1),(9,-2)\big)$
$(-2,\frac{1}{4},\frac{3}{2},\frac{4}{3})$	$\frac{1}{2}$	$\big(D,(2,1),(2,1)\big)\bigcup_{\begin{pmatrix}-1&4\\1&-3\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
$(-2,\frac{1}{3},\frac{2}{3},\frac{5}{3})$	$\frac{1}{2}$	$\big(S^2,(2,1),(3,-1),(5,2)\big)$

TABLE 4. α a filling instruction on M_5 not factoring through M_4 with $e\big(M_5(\alpha)\big)=4$, and $E_\tau\big(M_5(\alpha)\big)=\{\beta,0,1,\infty\}$, part 3/3.

α	Additional exceptional slopes β_i	Exceptional filling $M_4(\alpha)(\beta_i)$
$(-2,-2,-2)$	$\beta_1=-2$	$\big(D,(2,1),(2,-1)\big)\bigcup_{\begin{pmatrix}-1&4\\1&-3\end{pmatrix}}\big(D,(2,1),(3,1)\big)$
	$\beta_2=-1$	$T_{\begin{pmatrix}1-n&1\\-1&0\end{pmatrix}}$
$(-2,-\frac{1}{2},-2)$	$\beta_1=-1$	$\big(A,(2,1)\big)/_{\begin{pmatrix}-1&1\\1&0\end{pmatrix}}$
	$\beta_2=3$	$\big(D,(2,1),(2,-1)\big)\bigcup_{\begin{pmatrix}2&1\\-1&1\end{pmatrix}}\big(D,(2,1),(3,1)\big)$

TABLE 5. α a filling instruction on M_4 not factoring through M_3 with $e_\tau\big(M_4(\alpha)\big)=6$, and $E_\tau\big(M_4(\alpha)\big)=\{\beta_1,\beta_2,0,1,2,\infty\}$.

α	Additional exceptional slope β	Exceptional filling $M_4(\alpha)(\beta)$
$(-2, \frac{r}{s}, -2)$	$\beta = -1$	$\big(A, (s, s{-}r)\big) / \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
$(\frac{p}{q}, 4, -\frac{1}{2})$		$\big(D, (2,1), (q,p)\big) \bigcup_{\begin{pmatrix} -1 & 1 \\ 2 & -1 \end{pmatrix}} \big(D, (2,1), (3,1)\big)$
$(4, 5, -\frac{1}{2})$		$\big(A, (2,1)\big) / \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix}$
$(-2, 4, -\frac{2}{3})$		$\big(D, (2,1), (2,-1)\big) \bigcup_{\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}} \big(D, (2,1), (3,1)\big)$
$(-2, -5, -3)$		$\big(D, (2,1), (2,1)\big) \bigcup_{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}} \big(D, (2,1), (3,1)\big)$
$(-2, -2, -6)$		$\big(D, (2,1), (2,1)\big) \bigcup_{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}} \big(D, (2,1), (3,1)\big)$
$(-2, -2, -3)$		$\big(S^2, (2,-1), (3,1), (7,1)\big)$
$(-2, -3, -3)$,		$\big(S^2, (2,-1), (4,1), (5,1)\big)$
$(-2, -2, -4)$		$\big(S^2, (2,-1), (4,1), (5,1)\big)$
$(-3, -4, -2)$		$\big(S^2, (3,-2), (3,1), (4,1)\big)$
$(-2, -2, -5)$		$\big(S^2, (3,-2), (3,1), (4,1)\big)$
$(-3, -2, -3)$		$\big(D, (2,1), (2,-1)\big) \bigcup_{\begin{pmatrix} -1 & 3 \\ 1 & -2 \end{pmatrix}} \big(D, (2,1), (3,1)\big)$

TABLE 6. α a filling instruction on M_4 not factoring through M_3 with $e_\tau\big(M_4(\alpha)\big) = 5$, and $E_\tau\big(M_4(\alpha)\big) = \{\beta, 0, 1, 2, \infty\}$.

References

[Ago] Agol, I., *The minimal volume for orientable hyperbolic* 2*-cusped* 3*-manifolds* Proc. Amer. Math. Soc. **138** (2010), 3723-3732.

[Bak] Baker, K. L., *Surgery descriptions and volumes of Berge knots II: Descriptions of the Minimally Twisted Five Chain Link*, J. Knot Theory Ramifications, **17** (2008).

[CHW] Callahan, P.J., Hildebrand, M.V., Weeks, J.R., *A census of cusped hyperbolic* 3*-manifolds*, Math. Comp. **68** (1999), 321-332.

[GAS] González-Acuña, F., Short, H., *Knot surgery and primeness*, Math. Proc. Cambridge Philos. Soc. **99** (1986), 89-102.

[G1] Gordon, C., McA. *Dehn fillings: a survey*, in Knot Theory, Banach Center Publications **42**, Institute of Mathematics, Polish Academy of Sciences, Warsaw, 1998, 120-144.

[G2] Gordon, C., McA. *Dehn surgery and* 3*-manifolds*, Mrowka, T.S., Ozsváth (Eds.) IAS/Park City Mathematics Series, **16** (2006).

[GL1] Gordon, C. McA., Luecke, J., *Knots are determined by their complements* J. Amer. Math. Soc. **2** (1989), 371415.

[GL2] Gordon, C. McA., Luecke, J., *Dehn surgeries on knots creating essential tori, I*, Comm. Anal. Geom. **3** (1995), 371-644.

[Kan] Kang, S., *Reducible and toroidal Dehn fillings with distance* 3, Topology **47** (2008), 277-315.

[Kir] Kirby, K., *Problems in low-dimensional topology*, available from `http://math.berkeley.edu/~kirby/problems.ps`

[LM] Lackenby, M., Meyerhoff, R., *The maximal number of exceptional Dehn surgeries*, Invent. Math. (to appear), available at http://people.maths.ox.ac.uk/∼lackenby/ .

[Lee] Lee, S., *Lens spaces and toroidal Dehn fillings*, Math. Z. **267** (2011), 781-802.

[MP] Martelli, B., Petronio, C., *Dehn filling of the "magic"* 3*-manifold*, Comm. Anal. Geom. **14** (2006), 969-1026.

[MPR] Martelli, B., Petronio, C., Roukema, F.W.M., *Exceptional Dehn surgery on the minimally twisted* 5*-chain link*, available at `http://arxiv.org/PS_cache/arxiv/pdf/1109/1109.0903v1.pdf`

[Rol] Rolfsen, D., "Knots and Links", AMS Chelsea Publishing, (1976).

[Rou] Roukema, F.W.M., *Dehn surgery on the minimally twisted five-chain link*, Ph.D. thesis, supervised by C. Petronio and B. Martelli 2011, available at `http://etd.adm.unipi.it/theses/available/etd-09212011-102615/`

[Yos] Yoshida, K., *The minimal volume orientable hyperbolic 3-manifold with 4 cusps*, available at `http://arxiv.org/abs/1209.1374`.